

N°921 / PC

TOPIC(s) : Alternative technologies

Aqueous biphasic systems based on chiral ionic liquids as platforms for chiral resolution

AUTHORS

Tânia SINTRA / UNIVERSITY OF AVEIRO, CAMPUS UNIVERSITÁRIO DE SANTIAGO, AVEIRO

Mariam KHOLANY / UNIVERSITY OF AVEIRO, CAMPUS UNIVERSITÁRIO DE SANTIAGO, AVEIRO

Francisca SILVA / UNIVERSITY OF AVEIRO, CAMPUS UNIVERSITÁRIO DE SANTIAGO, AVEIRO

Sónia VENTURA / UNIVERSITY OF AVEIRO, CAMPUS UNIVERSITÁRIO DE SANTIAGO, AVEIRO

João COUTINHO / UNIVERSITY OF AVEIRO, CAMPUS UNIVERSITÁRIO DE SANTIAGO, AVEIRO

PURPOSE OF THE ABSTRACT

The production of enantiopure drugs is a major challenge for the pharmaceutical industry [1]. Considering the increased difficulty in the production of pure enantiomers by direct synthesis, the synthesis of racemates followed by their chiral resolution is a simpler, more flexible and cheaper alternative. This approach is commonly achieved by chromatography and crystallization, yet other methods, such as enantioselective liquid-liquid extraction can provide a better balance between costs, operational versatility and scale-up opportunities. Aqueous biphasic systems (ABS) are good candidates to turn enantioseparations not only into more biocompatible but also more versatile approaches [2]. Ionic liquids are alternative solvents with an enormous degree of structural diversity, allowing the design of task-specific solvents and, by their introduction in ABS, of highly performant extraction/separation approaches. Being made up of ions, if one can select/develop chiral structures to function as cations, anions or both, the opportunity of creating chiral ionic liquids (CILs) emerges [3,4].

This work proposes the development of a versatile enantioseparation platform for the chiral resolution of racemic mandelic acid. ABS using CILs simultaneously as chiral selectors and phase forming agents is proposed. Five different ABS were studied to determine the CIL structure and salt role on the ABS aptitude to separate mandelic acid enantiomers. Optimization studies were further pursued, as representative CIL-based ABS were employed in optimization studies, where the mandelic acid content, temperature, tie-line length, salt and phases weight ratio were studied. The influence of these parameters is shown to be highly dependent on the CIL-based ABS, nevertheless the results attained suggest that the key driving the enantioseparation in these ABS is a combination of the enantiorecognition ability of a given CIL with the solubility of mandelic acid in the corresponding CIL-rich phase. By closely manipulating the two conditions of the purification process, good enantioselective separations can be achieved.

Acknowledgements

This work was developed within the scope of the project CICECO-Aveiro Institute of Materials, FCT Ref. UID/CTM/50011/2019, financed by national funds through the FCT/MCTES. This work was also financially supported by the project POCI-01-0145-FEDER- (PIDC/EQU-EPQ/30750/2017)- funded by FEDER, through COMPETE2020 - Programa Operacional Competitividade e Internacionalização (POCI), and by national funds (OE), through FCT/MCTES. The authors are grateful for the financial support of the FCT for the doctoral grant of SFRH/BD/94901/2013 of F.A. e Silva, IF/00402/2015 of S.P.M. Ventura and for the project PAC ? Programa de atividades Conjuntas ref: SAICTPAC/0040/2015.

FIGURES

FIGURE 1

FIGURE 2

KEYWORDS

Aqueous biphasic systems | Chiral ionic liquids | Enantioseparation | Mandelic acid enantiomers

BIBLIOGRAPHY

- [1] S. Mane. *Analytical Methods*, 8(42) (2016) 7567-7586.
- [2] M.G Freire, A.F.M. Cláudio, J.M.M. Araújo, J.A.P. Coutinho, I.M. Marrucho, J.N.C. Lopes, L.P.N. Rebelo, *Chemical Society Reviews*, 41 (2012) 4966-4995.
- [3] H. Wu, S. Yao, G. Qian, T. Yao, H. Song. *Journal of Chromatography A*, 1418 (2015) 150-157.
- [4] D. Wu, Y. Zhou, P. Cai, S. Shen, Y. Pan. *Journal of Chromatography A*, 1395 (2015) 65-72.